

CLASSES OF BANACH SPACES STABLE AND UNSTABLE WITH RESPECT TO THE OPENING

M.I.Ostrovskii¹

Mathematics Department, Boğaziçi University
80815 Bebek, Istanbul, TURKEY

and

Mathematical Division, Institute for Low Temperature Physics
47 Lenin avenue, 310164 Kharkov, UKRAINE

e-mail: mostrovskii@ilt.kharkov.ua

Abstract. The paper is a complement to the survey: M.I.Ostrovskii "Topologies on the set of all subspaces of a Banach space and related questions of Banach space geometry", Quaestiones Math. (to appear). It contains proofs of some results on stability of properties of Banach spaces with respect to the geometric opening stated in the survey without proofs.

Some results of the present paper are of independent interest, in particular the description of a predual property of the Banach–Saks property.²

The present paper is a complement to the survey [O1]. It contains proofs of results stated in [O1] without proofs.

Let us recall some of the notions from [O1] and fix some notation. Let X be a Banach space. The closed unit ball and the unit sphere of a Banach space X are denoted by $B(X)$ and $S(X)$ respectively. Recall that the *density character* of a topological space X is defined to be the least cardinality of dense subset of X and is denoted by $\text{dens}X$. We shall denote the set of all closed subspaces of X by $G(X)$.

Suppose $Y, Z \in G(X)$. Let

$$\Theta_0(Y, Z) = \sup\{\text{dist}(y, Z) : y \in S(Y)\}$$

$$\Omega_0(Y, Z) = \sup\{\text{dist}(y, S(Z)) : y \in S(Y)\}.$$

It is clear that $0 \leq \Theta_0(Y, Z) \leq \Omega_0(Y, Z) \leq 2\Theta_0(Y, Z)$.

The *opening* between Y and Z is defined to be

$$\Theta(Y, Z) = \max\{\Theta_0(Y, Z), \Theta_0(Z, Y)\}.$$

¹Supported by TÜBİTAK

²1991 Mathematics Subject Classification 46B20

The *spherical opening* between Y and Z is defined to be

$$\Omega(Y, Z) = \max\{\Omega_0(Y, Z), \Omega_0(Z, Y)\}.$$

We shall not distinguish terms “class of Banach spaces” and “property of Banach spaces”.

Let P be a class of Banach spaces. Class P is called *stable* if there exists a number $\alpha > 0$ such that for every Banach space X and every $Y, Z \in G(X)$ the following implication holds

$$(Y \in P) \& (\Theta(Y, Z) < \alpha) \Rightarrow (Z \in P).$$

Class P is called *extendedly stable* if there exists a number $\alpha > 0$ such that for every Banach space X and every $Y, Z \in G(X)$ the following implication holds

$$(Y \in P) \& (\Theta_0(Z, Y) < \alpha) \Rightarrow (Z \in P).$$

One of the methods of establishing of unstability presented in [O1] is based on the following construction.

Let Y and Z be Banach spaces and let $T : S(Y) \rightarrow S(Z)$ and $D : S(Y^*) \rightarrow S(Z^*)$ be some surjective mappings (we suppose here that such mappings exist). Let us introduce on the algebraic sum $Y \oplus Z$ the following seminorm:

$$p(y, z) = \sup\{|y^*(y) - (Dy^*)(z)| : y^* \in S(Y^*)\}.$$

Seminorm p generates norm on the quotient of $Y \oplus Z$ by the zero-space of p . Denote the completion of this normed space by X .

By properties of D the spaces Y and Z are isometric to their natural images in X . It is not hard to verify (see [O1], 6.14) that the spherical opening between the images of Y and Z in X is not greater than the following value:

$$\sup\{|y^*(y) - (Dy^*)(Ty)| : y \in S(Y), y^* \in S(Y^*)\}. \quad (1)$$

Let us introduce the quantity $k(Y, Z)$ as the infimum of quantities (1) taken over all surjective mappings $T : S(Y) \rightarrow S(Z)$ and $D : S(Y^*) \rightarrow S(Z^*)$. If such mappings (at least one of them) do not exist we let $k(Y, Z) = 2$.

Let us observe that the value (1) is not greater than 2 for every Y, Z, T, D . So $k(Y, Z) \leq 2$ for every pair (Y, Z) .

Following M.I.Kadets [K] we introduce for every pair (Y, Z) of Banach spaces the value

$$d_\Omega(Y, Z) = \inf_{X, U, V} \Omega(UY, VZ)$$

where the infimum is taken over all Banach spaces X containing isometric copies of Y and Z and over all isometric embeddings $U : Y \rightarrow X$ and $V : Z \rightarrow X$.

Arguments above show that $d_\Omega(Y, Z) \leq k(Y, Z)$. In [O1] I asserted that these values are equivalent. Now we shall prove this assertion.

Proposition 1. *For every Banach spaces Y and Z the following inequality*

$$20d_\Omega(Y, Z) \geq k(Y, Z).$$

is valid.

Proof. From the inequality $k(Y, Z) \leq 2$ it follows that we may restrict ourselves to the case $d_\Omega(Y, Z) < 1/10$.

Let ε be an arbitrary number of the open interval $(0, 1)$. By definition of d_Ω there exists a Banach space X containing subspaces isometric to Y and Z (we shall still denote them by Y and Z), for which

$$\Omega(Y, Z) \leq d_\Omega(Y, Z)(1 + \varepsilon).$$

We shall use the following definition. Subset L of a metric space X is said to be τ -lattice ($\tau \in \mathbf{R}$, $\tau > 0$), if the distance between each two elements of L is not less than τ , and is said to be *maximal* τ -lattice, if L is not a proper subset of any τ -lattice in M . By standard application of Hausdorff's maximality theorem it follows that every metric space contains a maximal τ -lattice for arbitrary $\tau > 0$.

Let $\{y_\alpha\}_{\alpha \in A}$ be a maximal $(2(1+\varepsilon)\Omega(Y, Z))$ -lattice in $S(Y)$. By definition of maximal τ -lattice we can find such subsets $B_\alpha \subset S(Y)$ ($\alpha \in A$) that the following conditions are satisfied.

- (a) B_α contains the intersection of $S(Y)$ with the open ball of radius $(1 + \varepsilon)\Omega(Y, Z)$ centered at y_α .
- (b) B_α is contained in the ball of the radius $2(1 + \varepsilon)\Omega(Y, Z)$ centered at y_α .
- (c) Sets B_α are pairwise disjoint.
- (d) $\cup_{\alpha \in A} B_\alpha = S(Y)$.

For every $\alpha \in A$ we choose $z_\alpha \in S(Z)$ such that $\|z_\alpha - y_\alpha\| < (1 + \varepsilon)\Omega(Y, Z)$. It is easy to verify that $\{z_\alpha\}_{\alpha \in A}$ is a $4(1 + \varepsilon)\Omega(Y, Z)$ -net in $S(Z)$. Let us denote the intersection of $S(Z)$ with the ball of radius $4(1 + \varepsilon)\Omega(Y, Z)$ centered at z_α by R_α ($\alpha \in A$). We have $S(Z) = \cup_{\alpha \in A} R_\alpha$.

Since we may (and shall) assume without loss of generality that $\Omega(Y, Z) < 1/2$ (see remark at the begining of the proof), then $\text{dens} Y = \text{dens} Z$ (see [KKM] (Theorem 3) or [O1] (section 6.27.2)). Therefore by condition (a) $\text{card} R_\alpha = \text{card} B_\alpha$. So we can define a mapping $T : S(Y) \rightarrow S(Z)$ in such a way that $Ty_\alpha = z_\alpha$ and $T(B_\alpha) = R_\alpha$. It is clear that this mapping is surjective.

Let $y^* \in S(Y^*)$. It is easy to see that, if we extend it in a norm-preserving manner onto the whole X and then restrict this extension to Z , then the norm of the obtained functional is not less than $1 - \Omega_0(Y, Z)$.

Let $\{y_\gamma^*\}_{\gamma \in C}$ be some maximal $(2(1 + \varepsilon)\Omega(Y, Z))$ -lattice in $S(Y^*)$. It is easy to see that there exists subsets $B_\gamma^* \subset S(Y^*)$ ($\gamma \in C$) satisfying the following conditions:

- (a) B_γ^* contains the intersection of $S(Y^*)$ with the open ball of radius $(1 + \varepsilon)\Omega(Y, Z)$ centered at y_γ^* .
- (b) B_γ^* is contained in the ball of radius $2(1 + \varepsilon)\Omega(Y, Z)$ centered at y_γ^* .
- (c) Sets B_γ^* are pairwise disjoint.
- (d) $\cup_{\gamma \in C} B_\gamma^* = S(Y^*)$.

Let $\{f_\gamma\}_{\gamma \in C}$ be some norm-preserving extensions of functionals $\{y_\gamma^*\}_{\gamma \in C}$ onto X . Let $z_\gamma^* = f_\gamma|_Z / \|f_\gamma|_Z\|$. Show that $\{z_\gamma^*\}_{\gamma \in C}$ is an $(10(1 + \varepsilon)\Omega(Y, Z))$ -net in $S(Z^*)$.

Suppose the contrary. Then there exists $z^* \in S(Z^*)$ such that $\|z^* - z_\gamma^*\| > 10(1 + \varepsilon)\Omega(Y, Z)$ for every $\gamma \in C$.

Therefore for every $\gamma \in C$ there exists $z_\gamma \in S(Z)$ such that

$$|(z^* - f_\gamma / \|f_\gamma|_Z\|)(z_\gamma)| > 10(1 + \varepsilon)\Omega(Y, Z).$$

Let g be some norm-preserving extension of z^* onto X . Let $y_\gamma \in S(Y)$ ($\gamma \in C$) be such that $\|z_\gamma - y_\gamma\| < (1 + \varepsilon)\Omega(Y, Z)$. We have

$$|(g - f_\gamma / \|f_\gamma|_Z\|)(y_\gamma)| > 8(1 + \varepsilon)\Omega(Y, Z).$$

Therefore

$$|||f_\gamma|_Z|g|_Y - y_\gamma^*|| > (1 - \Omega(Y, Z))8(1 + \varepsilon)\Omega(Y, Z).$$

Since we assumed that $\Omega(Y, Z) < 1/2$, then

$$|||f_\gamma|_Z||g|_Y - y_\gamma^*|| > 4(1 + \varepsilon)\Omega(Y, Z).$$

Since the lattice $\{y_\gamma^*\}_{\gamma \in C}$ is maximal, it follows that for some $\gamma \in C$ we have

$$||g|_Y/||g|_Y|| - y_\gamma^*|| < 2(1 + \varepsilon)\Omega(Y, Z).$$

Hence

$$||y_\gamma^* - ||f_\gamma|_Z||g|_Y|| < 2(1 + \varepsilon)\Omega(Y, Z) + ||g|_Y|| (1/||g|_Y|| - ||f_\gamma|_Z||) < 4(1 + \varepsilon)\Omega(Y, Z).$$

This contradiction proves the assertion concerning $\{z_\gamma^*\}_{\gamma \in C}$. Let us denote the intersection of $S(Z^*)$ with the ball of radius $10(1 + \varepsilon)\Omega(Y, Z)$ with the centre in z_γ^* by R_γ^* ($\gamma \in C$). It is clear that $S(Z^*) = \cup_{\gamma \in C} R_\gamma^*$.

By $\Omega(Y, Z) < 1/2$ it follows (see e.g. [O2]) that $\text{dens} Y^* = \text{dens} Z^*$. Therefore for every $\gamma \in C$ we have $\text{card} R_\gamma^* = \text{card} B_\gamma^*$. Hence we can find a mapping $D : S(Y^*) \rightarrow S(Z^*)$ such that $D(y_\gamma^*) = z_\gamma^*$ and $D(B_\gamma^*) = R_\gamma^*$ for every $\gamma \in C$. It is clear that D is surjective.

Let us prove that the mappings D and T give us the desired estimate. Let $y^* \in B_\gamma^*$ and $y \in B_\alpha$. We have

$$\begin{aligned} & |y^*(y) - (Dy^*)(Ty)| \leq \\ & |y_\gamma^*(y_\alpha) - y_\gamma^*(y_\alpha - y) - (y_\gamma^* - y^*)y - (z_\gamma^*(z_\alpha) - z_\gamma^*(z_\alpha - Ty) - (z_\gamma^* - Dy^*)(Ty))| \leq \\ & |y_\gamma^*(y_\alpha) - z_\gamma^*(z_\alpha)| + 2(1 + \varepsilon)\Omega(Y, Z) + 2(1 + \varepsilon)\Omega(Y, Z) + \\ & 4(1 + \varepsilon)\Omega(Y, Z) + 10(1 + \varepsilon)\Omega(Y, Z) \leq \\ & |f_\gamma(y_\alpha - z_\alpha) - (f_\gamma - z_\gamma^*)(z_\alpha)| + 18(1 + \varepsilon)\Omega(Y, Z) \leq (1 + \varepsilon)\Omega(Y, Z) + \Omega(Y, Z) + \\ & 18(1 + \varepsilon)\Omega(Y, Z) \leq 20(1 + \varepsilon)\Omega(Y, Z). \end{aligned}$$

Hence

$$k(Y, Z) \leq 20(1 + \varepsilon)\Omega(Y, Z) \leq 20d_\Omega(Y, Z)(1 + \varepsilon)^2.$$

Since ε is arbitrary number from the open interval $(0, 1)$, we obtain the desired inequality. The proposition is proved. $\square$

The aim of the remaining part of the present paper is to support the conjecture on characterization of extendedly stable classes which was formulated in [O1] (section 6.36).

Recall necessary definitions.

Let Γ be a set, and let $l_1(\Gamma)$ be the corresponding space of functions on Γ . Let X be a Banach space. By $l_\infty(\Gamma, X)$ we denote the space of bounded functions $x : \Gamma \rightarrow X$ with the sup-norm.

Let A be a subset of the unit sphere of $l_1(\Gamma)$. For every $a \in A$ we introduce a linear operator from $l_\infty(\Gamma, X)$ into X defined by the rule

$$x \rightarrow \sum_{\gamma \in \Gamma} a(\gamma)x(\gamma).$$

This operator will be also denoted by a . It is clear that the norm of this operator equals 1. The supremum of those δ for which there exists $x \in S(l_\infty(\Gamma, X))$ such that $\inf_{a \in A} \|a(x)\| \geq \delta$ is called the *index of A in X* and is denoted by $h(X, A)$

It is known (see [O1] (section 6.27)) that for many common properties of Banach spaces (reflexivity, B -convexity, Banach-Saks property) sets Γ , $A \subset S(l_1(\Gamma))$ and $\tau > 0$ can be chosen in such a way that the absence of the property for a Banach space X is equivalent to each of the following inequalities:

$$h(X, A) > 0;$$

$$h(X, A) \geq \tau.$$

Properties admitting such descriptions are called *l_1 -properties*. It is not hard to verify that l_1 -properties are extendedly stable (see [O1], Proposition 6.21).

In [O1] I conjectured that the converse is also true, more precisely, that for every cardinal number α the intersection of every extendedly stable property with the set of Banach spaces with density character less than α is an l_1 -property.

We shall prove several results supporting this conjecture. In passing we shall find new classes of l_1 -properties.

We introduce a subclass of the class of all l_1 -properties and prove that all known at the moment extendedly stable classes belong to this subclass.

Definition 1. Class P of Banach spaces will be called a *regular l_1 -property* if there exist a real number $\delta > 0$, a set Γ and a subset $A \subset S(l_1(\Gamma))$ satisfying the conditions:

1. The set A consists of finitely non-zero vectors.
2. If $a_0 \in A$ then A contains all vectors $a \in S(l_1(\Gamma))$ for which

$$(\forall \gamma \in \Gamma)(\text{sign} a_0(\gamma) = \text{sign} a(\gamma)).$$

3. For a Banach space X the following conditions are equivalent:

- (a) $X \notin P$.
- (b) $h(X, A) > 0$.
- (c) $h(X, A) \geq \delta$.

Supremum of those $\delta > 0$ for which there exist Γ and $A \subset S(l_1(\Gamma))$ such that conditions 1, 2 and 3 are satisfied will be called the *regular exponent* of class P and will be denoted by $\text{reg}(P)$.

When working with regular l_1 -properties, we shall use the following definition and notation. Let $A \subset S(l_1(\Gamma))$. Mapping $\mathbf{s}: \Gamma \rightarrow \{0, 1, -1\}$ will be called an *admissible sign* for A if there exists $a \in A$ such that

$$(\forall \gamma \in \Gamma)(\mathbf{s}(\gamma) = \text{sign} a(\gamma)) \quad (2)$$

The set of all admissible signs for A will be denoted by $\mathbf{S}(A)$. Let $\mathbf{s}$ be an admissible sign for A . The set of all $a \in A$ for which (2) is satisfied will be denoted by $A(\mathbf{s})$. It is clear that $A = \cup_{\mathbf{s} \in \mathbf{S}(A)} A(\mathbf{s})$.

When working with regular l_1 -properties we shall repeatedly use the following lemma.

Lemma 1. *Let Γ be arbitrary set and let subset $A \subset S(l_1(\Gamma))$ be such that conditions 1 and 2 of Definition 1 are satisfied. Suppose the Banach space X and $x \in S(l_\infty(\Gamma, X))$ are such that*

$$\inf_{a \in A} \|a(x)\| \geq c. \quad (3)$$

Then for every $\mathbf{s} \in \mathbf{S}(A)$ there exists a functional $f_{\mathbf{s}} \in B(X^)$ such that for every $\gamma \in \text{supp } \mathbf{s}$ the following relations are valid:*

$$|f_{\mathbf{s}}(x(\gamma))| \geq c.$$

$$\text{sign} f_{\mathbf{s}}(x(\gamma)) = \mathbf{s}(\gamma).$$

Proof. Consider the set $\{\mathbf{s}(\gamma)x(\gamma) : \gamma \in \text{supp } \mathbf{s}\}$. By condition 2 of Definition 1 and inequality (3), the convex hull of this set does not intersect the open ball of radius c centered at zero. Using the separation theorem, we find required functional. $\square$

Now we are going to show that all l_1 -properties listed in [O1] (section 6.27) are regular l_1 -properties. Here I should note that at the moment I do not know l_1 -properties which are not regular.

1. Reflexivity. Description presented in [O1] (in 6.19) doesn't satisfy condition 2 of Definition 1. Nevertheless reflexivity is a regular l_1 -property. In order to prove this assertion we need another description of reflexivity (see [B1, p. 51]):

Let X be a Banach space. The following conditions are equivalent:

- (a) X is non-reflexive.
- (b) For every θ , $0 < \theta < 1$, there exist a sequence $\{x_k\}_{k \in \mathbf{N}}$ in $S(X)$ and a sequence $\{f_n\}_{n \in \mathbf{N}}$ in $S(X^*)$ such that:

$$f_n(x_k) = \begin{cases} \theta, & \text{if } n \leq k, \\ 0, & \text{if } n > k. \end{cases}$$

Let $A \subset S(l_1)$ be the set of all finitely non-zero vectors $a = \{a_k\}_{k=1}^\infty$ such that for some $n \in \mathbf{N}$ (depending on a) we have $a_k \leq 0$ when $k \geq n$ and $a_k \geq 0$ when $k < n$. It is clear that this set satisfies conditions 1 and 2 of Definition 1.

Let X be a Banach space. Let us prove that the following conditions are equivalent.

- (a) X is non-reflexive.
- (b) $h(X, A) > 0$.
- (c) $h(X, A) \geq 1/3$.

Since the set A contains the set considered in [O1] (6.19), we need to prove implication (a) $\Rightarrow$ (c) only.

Let $a \in A$ and let $n \in \mathbf{N}$ be such that $a_k \leq 0$ when $k \geq n$ and $a_k \geq 0$ when $k < n$. We have

$$f_1\left(\sum_{k=1}^{\infty} a_k x_k\right) = \theta\left(\sum_{k=1}^{n-1} |a_k| - \sum_{k=n}^{\infty} |a_k|\right).$$

$$f_n\left(\sum_{k=1}^{\infty} a_k x_k\right) = -\theta \sum_{k=n}^{\infty} |a_k|.$$

Hence

$$(f_1 - 2f_n)\left(\sum_{k=1}^{\infty} a_k x_k\right) = \theta \sum_{k=1}^{\infty} |a_k| = \theta.$$

Therefore $\|a(x)\| \geq \theta/3$. Since the description above is valid for every $\theta < 1$, it follows that $h(X, A) \geq 1/3$.

2. It is known (see [O1], Example 6.20) that for every uncountable cardinal α the class of all Banach spaces with density character less than α is an l_1 -property. But the sets introduced in [O1] do not satisfy Condition 2 of Definition 1. Nevertheless these properties are regular l_1 -properties. This can be shown in the following way. Let α be an uncountable cardinal and let Γ be a set of cardinality α . Let us introduce subset $A \subset S(l_1(\Gamma))$ as the set of all vectors with two-point support. It is clear that this set satisfies conditions 1 and 2 of Definition 1.

Let X be a Banach space. The following conditions are equivalent:

- (a) $\text{dens}X \geq \alpha$.
- (b) $h(X, A) > 0$.
- (c) $h(X, A) \geq 1/3$.

Since the introduced set contains one considered in [O1] (6.20), then we need to prove the implication (a) $\Rightarrow$ (c) only.

Let X be a Banach space with $\text{dens}X \geq \alpha$. We may consider Γ as a set of all ordinals which are less than the least ordinal of cardinality α . By well-known arguments (see [KKM, p. 98]) we can find subset $\{x(\gamma)\}_{\gamma \in \Gamma} \subset S(X)$ such that for every $\gamma \in \Gamma$ we have

$$\text{dist}(x(\gamma), \text{lin}\{x(\beta) : \beta < \gamma\}) > 1 - \varepsilon.$$

It is easy to verify that for every pair $\{\beta, \gamma\} \subset \Gamma$ with $\beta \neq \gamma$ we have

$$\inf\{\|rx(\gamma) + sx(\beta)\| : r, s \in \mathbf{R}, |r| + |s| = 1\} \geq (1 - \varepsilon)/(3 - \varepsilon).$$

Therefore for every $a \in A$ we have $\|a(x)\| \geq (1 - \varepsilon)/(3 - \varepsilon)$. Since ε is arbitrary, then $h(X, A) \geq 1/3$.

3. The class of finite dimensional Banach spaces. In this case we take $\Gamma = \mathbf{N}$ and $A \subset S(l_1)$ being the set of all vectors with two-point support. Using arguments from the previous example it is easy to verify that for a Banach space X the following statements are equivalent

- (a) X is infinite dimensional
- (b) $h(X, A) > 0$.
- (c) $h(X, A) \geq 1/3$.

Furthermore, using the main result of [P] it can be shown that the statements (a)–(c) are equivalent to $h(X, A) \geq 1/2$.

The sets introduced in [O1] (see 6.27) in order to describe classes 4–7 below satisfy all conditions of Definition 1. So these classes are regular l_1 -properties.

4. The class of Banach spaces whose dimension is not greater than n (for every $n \in \mathbf{N}$).

5. B -convexity.

6. The class of Banach spaces not containing isomorphic copies of l_1 .

7. Alternate signs Banach-Saks property.

8. Super-reflexivity. Subset of $S(l_1)$ introduced in [O1] to describe this property does not satisfy condition 2 of Definition 1. Therefore we introduce another subset $A \subset S(l_1)$. Let A be the set of all vectors of the following form:

$$(0, \dots, 0, a_1, \dots, a_n, 0, \dots),$$

where $\{a_i\}_{i=1}^n$ are such that for some $k \in \{1, \dots, n+1\}$ we have $a_i \geq 0$ when $i < k$ and $a_i \leq 0$ when $i \geq k$, and a_1 is preceded by $n(n-1)/2$ zeros.

It is clear that A satisfies conditions 1 and 2 of Definition 1.

Let X be a Banach space. Prove that the following conditions are equivalent:

(a) X is not super-reflexive.

(b) $h(X, A) > 0$.

(c) $h(X, A) = 1$.

Since the set A contains the set considered in [O1] the only thing which we need to prove is that (a) implies (c). In order to do this let us recall [B1, p. 270–271] that in every non-super-reflexive space for every $\varepsilon > 0$ and $n \in \mathbf{N}$ there exists a set $\{x_1, x_2, \dots, x_n\} \subset B(X)$ such that for every $1 \leq k \leq n$ we have

$$\|x_1 + \dots + x_k - x_{k+1} - \dots - x_n\| \geq n(1 - \varepsilon).$$

Let $\{a_i\}_{i=1}^n$ be any sequence satisfying the conditions above with $k \leq n$ (the case $k = n+1$ is easy). It is clear that

$$\left\| \sum_{i=1}^n a_i x_i \right\| \geq \|x_1 + \dots + x_k - x_{k+1} - \dots - x_n - \sum_{i=1}^k (1 - a_i)x_i + \sum_{i=k+1}^n (1 + a_i)x_i\| \geq$$

$$n(1 - \varepsilon) - \left(\sum_{i=1}^k |1 - a_i| + \sum_{i=k+1}^n |1 + a_i| \right) =$$

$$n(1 - \varepsilon) - \sum_{i=1}^k (1 - |a_i|) - \sum_{i=k+1}^n (1 - |a_i|) =$$

$$n(1 - \varepsilon) - (n - 1) = 1 - n\varepsilon.$$

The implication (a) $\Rightarrow$ (c) can be easily derived from this estimate.

9. Banach–Saks property.

Consider the set $A \subset S(l_1)$ consisting of all vectors satisfying the conditions

- (a) The cardinality of support is not greater than the least of its elements.
- (b) For every $\{a_k\}_{k=1}^\infty$ there exists $n \in \mathbf{N}$ such that $a_k \geq 0$ if $k < n$ and $a_k \leq 0$ if $k \geq n$.

B.Beauzamy [B2] proved that a Banach space has Banach–Saks property if and only if it is simultaneously reflexive and has alternate signs Banach–Saks property. Using this result, characterization of alternate signs Banach–Saks property (see [O1] (6.27.8)) and the above characterization of reflexivity it is not hard to verify that the following statements are equivalent:

- (a) X is not Banach–Saks
- (b) $h(X, A) > 0$.
- (c) $h(X, A) \geq 1/3$.

Proposition 2. *Let $\{P_\alpha\}_{\alpha \in C}$ be some set of regular l_1 -properties and the inequality*

$$\inf_{\alpha} \text{reg} P_\alpha > 0$$

is satisfied. Then $\cup_{\alpha \in C} P_\alpha$ is a regular l_1 -property and $\text{reg}(\cup_{\alpha \in C} P_\alpha) \geq \inf_{\alpha} \text{reg} P_\alpha$.

Proof. Let ε be an arbitrary number from the open interval $(0,1)$. Let Γ_α and $A_\alpha \subset S(l_1(\Gamma_\alpha))$ ($\alpha \in C$) be such that A_α satisfies conditions 1 and 2 of Definition 1 and for a Banach space X the following statements are equivalent:

- (a) $X \notin P_\alpha$.
- (b) $h(X, A_\alpha) > 0$.
- (c) $h(X, A_\alpha) > (1 - \varepsilon)\text{reg} P_\alpha$.

Let $\Gamma = \cup_{\alpha \in C} \Gamma_\alpha$. Let $A \subset S(l_1(\Gamma))$ be the union of natural images of A_α in $S(l_1(\Gamma))$. It is clear that A satisfies conditions 1 and 2 of Definition 1.

Let X be a Banach space. It is easy to see that the following statements are equivalent:

- (a) $X \notin (\cup_{\alpha \in C} P_\alpha)$.
- (b) $h(X, A) > 0$.
- (c) $h(X, A) \geq (1 - \varepsilon) \inf_{\alpha \in C} \text{reg} P_\alpha$.

Since $\varepsilon \in (0,1)$ is arbitrary, the proposition is proved. $\square$

Definition 2. Let P be some class of Banach spaces. *Preclass* of P is defined to be the class of all Banach spaces whose duals belong to P . Preclass

of P is denoted by $\text{pre}(P)$.

It is known that if P is extendedly stable then $\text{pre}(P)$ is also extendedly stable (for different versions and proofs of this result see [J], [AAG] and [O1] (Proposition 6.34)).

Theorem 1. *If P is a regular l_1 -property then $\text{pre}(P)$ is also a regular l_1 -property and $\text{reg}(\text{pre}(P)) \geq \text{reg}(P)$.*

Proof. Let θ be an arbitrary number from the open interval $(0,1)$. Let us choose a set Γ and a subset $A \subset S(l_1(\Gamma))$ in such a way that all conditions of Definition 1 with $\delta = \theta \text{reg}(P)$ are satisfied. Put $\Psi = \mathbf{S}(A)$. Let $D \subset S(l_1(\Psi))$ be a subset consisting of all finitely non-zero vectors d satisfying the following condition. If $\text{supp} d = \{\mathbf{s}_1, \dots, \mathbf{s}_n\}$, then for some

$$\gamma \in \cap_{i=1}^n \text{supp } \mathbf{s}_n$$

we have

$$(\forall i \in \{1, \dots, n\})(\text{sign} d(\mathbf{s}_i) = \mathbf{s}_i(\gamma)).$$

It is clear that D satisfies the conditions 1 and 2 of Definition 1. In order to prove Theorem 1 it is sufficient to prove that the following statements are equivalent.

- (a) $X \notin \text{pre}(P)$.
- (b) $h(X, D) > 0$.
- (c) $h(X, D) \geq \theta \text{reg}(P)$.

It is clear that we need to prove implications (a) $\Rightarrow$ (c) and (b) $\Rightarrow$ (a) only.

Let δ be an arbitrary number from the open interval $(0,1)$. Let $X \notin \text{pre}(P)$. It means that $X^* \notin P$. Let vector $x^* \in S(l_\infty(\Gamma, X^*))$ be such that

$$\inf_{a \in A} \|a(x^*)\| > \theta \delta \text{reg} P.$$

By Lemma 1 we can find for every $\mathbf{s} \in \mathbf{S}(A)$ a functional $f_{\mathbf{s}} \in B(X^{**})$ such that for every $\gamma \in \text{supp } \mathbf{s}$ we have $|f_{\mathbf{s}}(x^*(\gamma))| > \delta \theta \text{reg}(P)$ and $\text{sign} f_{\mathbf{s}}(x^*(\gamma)) = \mathbf{s}(\gamma)$. It is clear that we may suppose that $\|f_{\mathbf{s}}\| < 1$.

Further we shall repeatedly use the following statement which goes back to E.Helly (see [B1, p. 52]). Let $f \in X^{**}$ and $\|f\| < 1$. Let $\{x_1^*, x_2^*, \dots, x_n^*\}$ be a finite subset of X^* . Then there exists a vector $x \in X$ such that $\|x\| < 1$ and $x_i^*(x) = f(x_i^*)$ for every $i \in \{1, \dots, n\}$. Every such vector we shall call a *reflection of f with respect to $\{x_1^*, x_2^*, \dots, x_n^*\}$* .

Let $\{x(\mathbf{s})\}_{\mathbf{s} \in \Psi}$ be some reflections of $\{f_{\mathbf{s}}\}_{\mathbf{s} \in \Psi}$ with respect to sets

$$\{x^*(\gamma)\}_{\gamma \in \text{supp } \mathbf{s}}.$$

Let $d \in D$. Let us estimate from below the value $\|d(x)\|$. Let $\text{supp } d = \{\mathbf{s}_1, \mathbf{s}_2, \dots, \mathbf{s}_n\}$. Let $\gamma \in \cap_{i=1}^n \text{supp } \mathbf{s}_i$ be such that $\text{sign } d(\mathbf{s}_i) = \mathbf{s}_i(\gamma)$ for $i \in \{1, \dots, n\}$. We have

$$\begin{aligned} \|d(x)\| &= \left\| \sum_{i=1}^n d(\mathbf{s}_i)x(\mathbf{s}_i) \right\| \geq x^*(\gamma) \left(\sum_{i=1}^n d(\mathbf{s}_i)x(\mathbf{s}_i) \right) = \\ &= \sum_{i=1}^n d(\mathbf{s}_i)f_{\mathbf{s}_i}(x^*(\gamma)) = \sum_{i=1}^n \mathbf{s}_i(\gamma)|d(\mathbf{s}_i)|\mathbf{s}_i(\gamma)|f_{\mathbf{s}_i}(x^*(\gamma))| \geq \delta \theta \text{reg}(P). \end{aligned}$$

Since δ is arbitrary number from the open interval $(0,1)$, the argument above implies that (a) $\Rightarrow$ (c).

Let us turn to the implication (b) $\Rightarrow$ (a). Let $h(X, D) > \varepsilon > 0$. By Lemma 1 and weak* compactness of $B(X^*)$, for every $\gamma \in \Gamma$ we can find $x^*(\gamma) \in B(X^*)$ such that $|x^*(\gamma)x(\mathbf{s})| > \varepsilon$ and $\text{sign}(x^*(\gamma)x(\mathbf{s})) = \mathbf{s}(\gamma)$ for every $\mathbf{s}$ for which $\gamma \in \text{supp } \mathbf{s}$. Let $a \in A(\mathbf{s})$. We have

$$\left\| \sum_{\gamma \in \Gamma} a(\gamma)x^*(\gamma) \right\| \geq \left(\sum_{\gamma \in \text{supp } \mathbf{s}} a(\gamma)x^*(\gamma) \right)x(\mathbf{s}) = \sum_{\gamma \in \Gamma} |a(\gamma)|\mathbf{s}(\gamma)|x^*(\gamma)x(\mathbf{s})|\mathbf{s}(\gamma) > \varepsilon.$$

Hence $X^* \notin P$ and $X \notin \text{pre}(P)$. Theorem 1 is proved. $\square$

Remark. If property P has a concrete description as a regular l_1 -property, then the proof of Theorem 1 gives us a concrete description of $\text{pre}(P)$. Hence we obtain e.g. a description of pre-Banach-Saks property.

Definition 3. Let P be a class of Banach spaces. *Coclass* of P is defined to be the class of all Banach spaces for which the quotient space X^{**}/X is in P . Coclass of P is denoted by P^{co} .

J.Alvarez, T.Alvarez and M.Gonzalez [AAG] proved that the extended stability of P implies the extended stability of P^{co} . Our aim is to prove the following result.

Theorem 2. *Let P be a regular l_1 -property. Then P^{co} is also a regular l_1 -property.*

Proof. Let a set Γ , a subset $A \subset S(l_1(\Gamma))$ and a number $\delta > 0$ be such that all conditions of Definition 1 are satisfied. Let Ψ be the set of all triples (γ, α, k) , where $\gamma \in \Gamma$, α is a finite subset of $\mathbf{S}(A)$, $k \in \mathbf{N}$.

We define subset $E \subset S(l_1(\Psi))$ in the following way. Let $\{\alpha_1, \dots, \alpha_r\}$ be a finite collection of finite subsets of $\mathbf{S}(A)$ such that the following two conditions are satisfied:

$$\cap_{k=1}^r \alpha_k \neq \emptyset.$$

For some $m \in \{1, \dots, r-1\}$ and some finite subset $\alpha \subset \mathbf{S}(A)$ we have $\alpha \supset \alpha_k$ for $k \in \{1, \dots, m\}$ and $\alpha \subset \alpha_k$ for $k \in \{m+1, \dots, r\}$.

Let $n \in \mathbf{N}$ and $\mathbf{s} \in \cap_{k=1}^r \alpha_k$. We define $E(\{\alpha_1, \dots, \alpha_r\}, \mathbf{s}, n, m)$ as the set of finitely non-zero vectors of $S(l_1(\Psi))$ satisfying the following three conditions.

(a) Their supports are contained in the union of the following two sets:

$$\text{supps} \times \{\alpha_1, \dots, \alpha_m\} \times \{1, \dots, n\},$$

$$\text{supps} \times \{\alpha_{m+1}, \dots, \alpha_r\} \times \{n+1, \dots\}.$$

(b) The signs of all non-zero coordinates corresponding to the triples (γ, α_k, j) of the first set are equal to $\mathbf{s}(\gamma)$.

(c) The signs of all non-zero coordinates corresponding to the triples (γ, α_k, j) of the second set are equal to $-\mathbf{s}(\gamma)$.

The union of all sets $E(\{\alpha_1, \dots, \alpha_r\}, \mathbf{s}, n, m)$ constructed in the described way we shall denote by E . It is clear that E satisfies Conditions 1 and 2 of Definition 1.

Let X be a Banach space. Let us show that the following statements are equivalent.

(a) $X \notin P^{\text{co}}$.

(b) $h(X, E) > 0$.

(c) $h(X, E) \geq \delta/3$.

It is clear that we need to prove the implications (a) $\Rightarrow$ (c) and (b) $\Rightarrow$ (a) only.

Let us start with the implication (a) $\Rightarrow$ (c). Let $0 < \theta < 1$. By Lemma 1 we can find $u \in S(l_\infty(\Gamma, X^{**}/X))$ such that for every $\mathbf{s} \in \mathbf{S}(A)$ there exists $f_{\mathbf{s}} \in (X^{**}/X)^* = X^\perp \subset X^{***}$ such that $\|f_{\mathbf{s}}\| < 1$ and for every $\gamma \in \text{supps}$ we have $|f_{\mathbf{s}}(u(\gamma))| > \delta\theta$ and $\text{sign} f_{\mathbf{s}}(u(\gamma)) = \mathbf{s}(\gamma)$. It is clear that all conditions above are satisfied for some $u \in l_\infty(\Gamma, X^{**}/X)$ satisfying the inequality $\|u\| < 1$. So we shall suppose that $\|u\| < 1$.

Let $x^{**}(\gamma) \in X^{**}$ ($\gamma \in \Gamma$) be such that the image of $x^{**}(\gamma)$ under the quotient mapping $X^{**} \rightarrow X^{**}/X$ coincides with $u(\gamma)$ and $\|x^{**}(\gamma)\| < 1$.

Now we define $x \in S(l_\infty(\Psi, X))$ for which $\inf_{e \in E} \|e(x)\|$ is “large”. This function will be defined in stages. At first we define this function for triples (γ, α, k) with one-element set α , then for triples with two-element set α , etc.

It is clear that it is sufficient to define $x(\gamma, \alpha, k)$ only for those triples for which $\gamma \in \cup_{\mathbf{s} \in \alpha} \text{supps}$. (Since for other triples (γ, α, k) the numbers $e(\gamma, \alpha, k)$ are equal to zero for every $e \in E$ and so $x(\gamma, \alpha, k)$ can be defined arbitrarily.)

In the course of construction we shall also define functionals $f(\mathbf{s}, \alpha, k)$, where α is a finite subset of $\mathbf{S}(A)$, $\mathbf{s} \in \alpha$ and $k \in \mathbf{N}$.

For one-element sets α we define $x(\gamma, \alpha, k)$ by induction on k .

1. Let $\alpha = \{\mathbf{s}\}$.

1.1. Define $f(\mathbf{s}, \{\mathbf{s}\}, 1)$ as a reflection of $f_{\mathbf{s}}$ with respect to $\{x^{**}(\gamma) : \gamma \in \text{supps}\}$. Let $x(\gamma, \{\mathbf{s}\}, 1)$ be reflections of $\{x^{**}(\gamma) : \gamma \in \text{supps}\}$ with respect to $\{f(\mathbf{s}, \{\mathbf{s}\}, 1)\}$.

...

1.k. Define $f(\mathbf{s}, \{\mathbf{s}\}, k)$ as a reflection of $f_{\mathbf{s}}$ with respect to

$$\{x^{**}(\gamma) : \gamma \in \text{supps}\} \cup \{x(\gamma, \{\mathbf{s}\}, j) : \gamma \in \text{supps}, j = 1, \dots, k-1\}.$$

Let $x(\gamma, \{\mathbf{s}\}, k)$ be reflections of $\{x^{**}(\gamma) : \gamma \in \text{supps}\}$ with respect to

$$\{f(\mathbf{s}, \{\mathbf{s}\}, j) : j = 1, \dots, k\}.$$

...

Let us suppose that we have already constructed $x(\gamma, \alpha, k)$ for all $(m-1)$ -element subsets $\alpha \subset \mathbf{S}(A)$.

m. Let $\alpha = \{\mathbf{s}_1, \dots, \mathbf{s}_m\}$. By $\text{supp}\alpha$ we shall denote $\cup_{i=1}^m \text{supps}_{\mathbf{s}_i}$.

m.1. Define $\{f(\mathbf{s}_i, \alpha, 1)\}$ as a reflections of $f_{\mathbf{s}_i}$ with respect to $\{x^{**}(\gamma) : \gamma \in \text{supp}\alpha\}$. Let $\{x(\gamma, \alpha, 1) : \gamma \in \text{supp}\alpha\}$ be reflections of $\{x^{**}(\gamma) : \gamma \in \text{supp}\alpha\}$ with respect to $\{f(\mathbf{s}, \beta, 1) : \beta \subset \alpha, \mathbf{s} \in \beta\}$.

...

m.k. Define $\{f(\mathbf{s}_i, \alpha, k)\}$ as a reflections of $f_{\mathbf{s}_i}$ with respect to

$$\{x^{**}(\gamma) : \gamma \in \text{supp}\alpha\} \cup \{x(\gamma, \beta, j) : \beta \subset \alpha, \gamma \in \text{supp}\beta, j = 1, \dots, k-1\}.$$

Let $\{x(\gamma, \alpha, k)\}$ be reflections of $\{x^{**}(\gamma) : \gamma \in \text{supp}\alpha\}$ with respect to $\{f(\mathbf{s}, \beta, j) : \beta \subset \alpha, \mathbf{s} \in \beta, j = 1, \dots, k\}$.

...

Let us show that

$$(\forall e \in E)(\|e(x)\| \geq \delta\theta/3).$$

Let $e \in E(\{\alpha_1, \dots, \alpha_r\}, \mathbf{s}, n, m)$, where $\mathbf{s} \in \cap_{i=1}^r \alpha_i$. We have

$$e(x) = \sum_{\gamma \in \text{supps}} \left(\sum_{i=1}^m \sum_{k=1}^n e(\gamma, \alpha_i, k) x(\gamma, \alpha_i, k) + \sum_{i=m+1}^r \sum_{k=n+1}^{\infty} e(\gamma, \alpha_i, k) x(\gamma, \alpha_i, k) \right).$$

Therefore we have

$$\begin{aligned} f(\mathbf{s}, \{\mathbf{s}\}, 1)(e(x)) &= \sum_{\gamma \in \text{supps}} \left(\sum_{i=1}^m \sum_{k=1}^n e(\gamma, \alpha_i, k) x^{**}(\gamma) f(\mathbf{s}, \{\mathbf{s}\}, 1) + \right. \\ &\quad \left. \sum_{i=m+1}^r \sum_{k=n+1}^{\infty} e(\gamma, \alpha_i, k) x^{**}(\gamma) f(\mathbf{s}, \{\mathbf{s}\}, 1) \right) = \\ &= \sum_{\gamma \in \text{supps}} \left(\sum_{i=1}^m \sum_{k=1}^n |e(\gamma, \alpha_i, k)| |f_{\mathbf{s}}(x^{**}(\gamma))| - \sum_{i=m+1}^r \sum_{k=n+1}^{\infty} |e(\gamma, \alpha_i, k)| |f_{\mathbf{s}}(x^{**}(\gamma))| \right) \end{aligned}$$

According to definition of m there exists $\alpha \subset \mathbf{S}(A)$ such that $\cup_{i=1}^m \alpha_i \subset \alpha$ and $\alpha \subset \alpha_i$ when $i \geq m+1$. We have

$$\begin{aligned} f(\mathbf{s}, \alpha, n+1)(e(x)) &= \sum_{\gamma \in \text{supps}} \left(\sum_{i=1}^m \sum_{k=1}^n e(\gamma, \alpha_i, k) f(\mathbf{s}, \alpha, n+1) x(\gamma, \alpha_i, k) + \right. \\ &\quad \left. \sum_{i=m+1}^r \sum_{k=n+1}^{\infty} e(\gamma, \alpha_i, k) f(\mathbf{s}, \alpha, n+1) x(\gamma, \alpha_i, k) \right). \end{aligned}$$

Since $\alpha_i \subset \alpha$ if $i \leq m$ and $f_{\mathbf{s}} \in X^{\perp}$ then the first sum equals zero. Since $\alpha \subset \alpha_i$ when $i > m$, the second sum can be rewritten in the following way:

$$\sum_{m+1}^r \sum_{k=n+1}^{\infty} e(\gamma, \alpha_i, k) x^{**}(\gamma) f(\mathbf{s}, \alpha, n+1) = - \sum_{i=m+1}^r \sum_{k=n+1}^{\infty} |e(\gamma, \alpha_i, k)| |f_{\mathbf{s}}(x^{**}(\gamma))|.$$

Hence

$$\begin{aligned} 2|f(\mathbf{s}, \alpha, n+1)(e(x))| + f(\mathbf{s}, \{\mathbf{s}\}, 1)(e(x)) &= \\ \sum_{\gamma} \sum_i \sum_k |e(\gamma, \alpha_i, k)| |f_{\mathbf{s}}(x^{**}(\gamma))| &> \delta\theta. \end{aligned}$$

Therefore $\|e(x)\| \geq \delta\theta/3$. Since θ is arbitrary, the implication (a) $\Rightarrow$ (c) is proved.

Now we turn to the implication (b) $\Rightarrow$ (a). Let $h(X, E) = \tau > 0$. Let θ be arbitrary number from the open interval $(0, 1)$ and let $x \in S(l_{\infty}(\Psi, X))$ be such that $\inf_{e \in E} \|e(x)\| > \tau\theta$.

The set of pairs (α, k) , where α is a finite subset of $\mathbf{S}(A)$ and $k \in \mathbf{N}$ is a directed set with respect to the natural order. Let U be some ultrafilter which majorize the filter induced by this order. Let

$$x^{**}(\gamma) = w^* - \lim_U x(\gamma, \alpha, k) \quad (\gamma \in \Gamma).$$

Thus we defined $x^{**} \in B(l_\infty(\Gamma, X^{**}))$. It is clear that in order to finish the proof of Theorem 2 it is sufficient to prove that

$$(\forall a \in A)(\|\varphi \sum_{\gamma} a(\gamma) x^{**}(\gamma)\| \geq \tau\theta/2,$$

where φ is the quotient mapping $\varphi : X^{**} \rightarrow X^{**}/X$.

Assume the contrary, i.e. there exists $a \in A$ and $x \in X$ such that

$$\|\sum_{\gamma} a(\gamma) x^{**}(\gamma) - x\| < \tau\theta/2.$$

By the definition of $x^{**}(\gamma)$ for every finite subset $\alpha_0 \subset \mathbf{S}(A)$ and every $k_0 \in \mathbf{N}$ we have

$$\sum_{\gamma} a(\gamma) x^{**}(\gamma) \in w^* - \text{cl}\{\sum_{\gamma} a(\gamma) x(\gamma, \alpha, k) : \alpha \supset \alpha_0, k \geq k_0\}.$$

By Proposition 6.30 of [O1] we can find a collection $\{\alpha_1, \dots, \alpha_m\}$ of finite subsets of $\mathbf{S}(A)$, real numbers $\tau_1, \dots, \tau_m \geq 0$ with $\sum_{i=1}^m \tau_i = 1$ and numbers $k_1, \dots, k_m \in \mathbf{N}$ such that $\text{sign } a \in \cap_{i=1}^m \alpha_m$ and

$$\|\sum_{\gamma} a(\gamma) x^{**}(\gamma) - \sum_{i=1}^m \tau_i \sum_{\gamma} a(\gamma) x(\gamma, \alpha_i, k_i)\| < \tau\theta.$$

Using Proposition 6.30 of [O1] once more we find a collection $\{\alpha_{m+1}, \dots, \alpha_r\}$ of finite subsets of $\mathbf{S}(A)$, satisfying the condition $\cap_{i=m+1}^r \alpha_i \supset \cup_{i=1}^m \alpha_i$, real numbers $\tau_{m+1}, \dots, \tau_r \geq 0$ with $\sum_{i=m+1}^r \tau_i = 1$ and numbers $k_{m+1}, \dots, k_r \in \mathbf{N}$ such that

$$\max_{1 \leq i \leq m} k_i < \min_{m+1 \leq i \leq r} k_i$$

and

$$\|\sum_{\gamma} a(\gamma) x^{**}(\gamma) - \sum_{i=m+1}^r \tau_i \sum_{\gamma} a(\gamma) x(\gamma, \alpha_i, k_i)\| < \tau\theta.$$

Hence

$$\left\| \sum_{i=1}^m \tau_i \sum_{\gamma} a(\gamma) x(\gamma, \alpha_i, k_i) - \sum_{i=m+1}^r \tau_i \sum_{\gamma} a(\gamma) x(\gamma, \alpha_i, k_i) \right\| < 2\tau\theta.$$

It remains to note that the vector in the left-hand side of this inequality has the form $2e(x)$ for some $e \in E$. We arrived at a contradiction. The theorem is proved. $\square$

References

- [AAG] J.Alvarez, T.Alvarez and M.Gonzalez, The gap between subspaces and perturbation of non semi-Fredholm operators, Bull. Austral. Math. Soc. 45 (1992) no. 3, pp. 369–376
- [B1] B.Beauzamy, Introduction to Banach Spaces and their Geometry, Amsterdam, North-Holland Publishing Company, 1982
- [B2] B.Beauzamy, Banach-Saks properties and spreading models, Math. Scand. 44 (1979) no. 2, pp. 357–384
- [J] R.Janz, Pertubation of Banach spaces, preprint, Konstanz, 1987
- [K] M.I.Kadets, Note on the gap between subspaces (Russian), Funkts. Anal. Prilozhen. 9 (1975) no. 2, pp. 73–74
- [KKM] M.G.Krein, M.A.Krasnoselskii and D.P.Milman, On the defect numbers of linear operators in a Banach space and on some geometric questions (Russian), Sbornik Trudov Inst. Matem. AN Ukrainian SSR 11 (1948) pp. 97–112
- [O1] M.I.Ostrovskii, Topologies on the set of all subspaces of a Banach space and related questions of Banach space geometry, Quaestiones Math. (to appear)
- [O2] M.I.Ostrovskii, The relation of the gaps of subspaces to dimension and other isomorphic invariants (Russian), Akad. Nauk Armyan. SSR Dokl. 79 (1984) no. 2 pp. 51–53
- [P] A.Pelczyński All separable Banach spaces admit for every $\varepsilon > 0$ fundamental total and bounded by $1 + \varepsilon$ biorthogonal sequences, Studia Math. 55 (1976) pp. 295–304